



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Climatology Ocean Emissivity Trends 2010-04-23

Evan Manning

Jet Propulsion Laboratory, California Institute of Technology

Copyright 2010
California Institute of Technology
Government sponsorship acknowledged

ECMWF Climatology
AIRS Science Team Meeting
April 21-23 2010, Pasadena CA



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Climatology

- **I Discussed ECMWF Climatology in May & Sept 2009 AIRS Science Team meetings**
 - **We extract ECMWF forecast at AIRS overpass locations/times**
 - Separate by land/ocean
 - Produce Level-2 simulated files
 - **Process to Level-3 daily and monthly files**
 - **Roll up multiple years to monthly climatology files in AIRS Level-3 format**
 - **Software used in Level-2 retrieval extracts appropriate info given month, latitude, longitude, ascending vs. descending, land fraction**
 - Files and software are available to the science team



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Climatology

- **Progress since September 2009:**
 - **ECMWF climatology now replaces the old NCEP/UARS climatology as the default background in the JPL retrieval system**
 - This has limited effect on retrievals because both MW-only and regression completely ignore the background except above 100 hPa
 - When both regression and MW-only retrieval steps are skipped this leaves a better starting point for cloud clearing + physical retrieval
 - **More ECMWF forecast data has been received.**
 - 6+ years with many small gaps and a few large ones.
 - Thanks Scott!
 - **More ECMWF forecast has been processed to L3 for incorporation.**
 - 59 months 2003-2009
 - **New ECMWF Clim with 3-4 years for each month (2005-8)**
 - Soon to be 4-6 with the addition of 2009
 - Standard deviation now more fully reflects interannual variability




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

ECMWF Climatology Gotchas: Tropopause

Atmospheric Infrared Sounder



- This is a histogram of the tropopause pressure
- Before Feb 2006 ECMWF had 60 vertical layers
- Since Feb 2006 ECMWF has 91 vertical layers
 - Spatial resolution also increased from $\frac{1}{2}$ to $\frac{1}{4}$ degree
- Interpolation of temperature profiles gives some layers which cannot have a tropopause or other kink
- This effect can be seen in AIRS v5 retrievals because pre-2006 ECMWF was used to train the regression
 - If we're going to use tropopause-relative trace-gas climatologies then we really need a good tropopause



National Aeronautics and
Space Administration
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Climatology Gotchas: Tropical Tropopause Detail



- **Since February 2006 (91 layers):**
 - The distribution of tropopause is relatively continuous.
 - The peak of the distribution of tropical tropopause is ~93 hPa.
- **Before February 2006 (60 layers):**
 - The distribution is discontinuous at bin boundaries
 - The peak is sharper and is shifted to ~100 hPa



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Gotchas: ECMWF changes

- **ECMWF is an operational forecast, not a consistent reanalysis.**
- **They update their model frequently**
 - **http://www.ecmwf.int/products/data/operational_system/evolution/**




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

ECMWF Climatology Gotchas: Day/Night Tsurf differences

Atmospheric Infrared Sounder



- Each quadrant shows a time series of day – night surface temperature 2005-2009
- For land, AIRS and ECMWF agree well
- For ocean, ECMWF showed no day/night differences until Dec 2008

Copyright 2010
California Institute of Technology
Government sponsorship acknowledged


ECMWF Climatology
AIRS Science Team Meeting
April 21-23 2010, Pasadena CA




National Aeronautics and
Space Administration
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

ECMWF Climatology Gotchas: Day/Night Tsurf differences

Atmospheric Infrared Sounder



- **ECMWF Ocean showed no day/night differences until Dec 2008**
- **Zonal means now vary seasonally between ~0.0 and ~0.5 K.**
- **Still somewhat smaller than AIRS**




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Gotchas: Stratospheric H₂O



- The AIRS L2 retrieval largely uses its climatology guess H₂O above 100 hPa unchanged
 - It is adjusted to match at 100 hPa
- For this tropical case:
 - The new ECMWF clim is drier than the old NCEP/UARS clim
 - But the shape is similar so, after shifting, the retrieved strat H₂O is similar




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ECMWF Gotchas: Stratospheric H₂O



- For this south polar case:
 - The new ECMWF clim is much drier than the old NCEP/UARS clim
 - The shape is very different around 100 hPa.
 - The retrieved strat H₂O is much drier



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Climatology Conclusions

Problems arise with a simple collection of ECMWF data for climatology:

- **Bad tropopause structure if we use data before February 2006**
- **Problems with polar stratospheric H₂O**
- **Day/Night differences underpredicted if we use data before December 2009**

An excellent AIRS-specific climatology could be built if demand is high enough:

- **Adjust polar 100 hPa structure**
- **Merge different fields from different time ranges**
 - **Standard deviations from the entire mission to get interannual variability**
 - **Fine vertical structure using one data since Feb 2006**
 - **Ocean Tsurf only since Dec 2009**



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Bonus Topic: Ocean Emissivity Trends


V5.0 vs. V5.6



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropical ocean emissivity spectrum v5.0




- **V5.0 tropical ocean emissivity:**
 - Had a clear increasing trend at all frequencies
 - Had large day/night differences in the shortwave region



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropical ocean emissivity spectrum v5.6




- **V5.6 tropical ocean emissivity:**
 - **Decreases trends at all frequencies**
 - Still has residual trends 1200-1400 & 2200-2400 cm⁻¹
 - These are not window regions
 - Trend in 2200-2400 cm⁻¹ may be only in daytime
 - **Decreases day/night differences in the shortwave region**
 - **Introduces a new kink in daytime shortwave emissivity**




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropical ocean emissivity spectra






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Ocean Emissivity by Month



- **V5.0 emissivity had a strong seasonal cycle in nighttime emissivity**
- **V5.6 does not**



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Emissivity Conclusions

- **V5.6 ocean emissivity is better by most metrics:**
 - ✓ **Trend reduction**
 - X But some residual trend remains
 - ✓ **Day vs. night differences**
 - ✓ **Seasonal cycle**
 - ✓ **All cases with Emissivity > 1.0 eliminated**
 - ✓ **Cases with Emissivity < 0.95 greatly reduced**
 - X **Shortwave structure introduced**